





















- Barker E. Neuroscience Nursing: A spectrum of care. 3<sup>rd</sup> ed. Missouri: Mosby Elsevier. 2009.
- Carroll LJ, Cassidy DJ, Peloso MP, Borg J, Holst VH, Holm L, Paniak C, & Pepin M. Prognosis for mild traumatic brain injury: Results of the WHO collaborating centre task force on mild traumatic brain injury. *Journal of Rehabilitation Medicine Supplement* 2004; 43: 84-105.
- Emanuelson I, Anderson E, & Holmkvist. Quality of life and post concussion symptoms in Adults after mild traumatic brain injury: a population-based study in western Sweden. *Acta Neurologica Scandinavica* 2003; 108: 332-338.
- King NS, Crawford S, Wenden FJ, Moss NG, & Wade DT. The Rivermead Post Concussion Symptoms Questionnaire: A measure of symptoms commonly experienced after head injury and its reliability. *Journal of Neurology* 1995; 242 (9): 587-592.
- Leininger BE, Gramling SE, Farrell AD, Kreutzer JS, & Peck EA. Neuropsychological deficits in symptomatic minor head injury patients after concussion and mild concussion. *Journal of Neurology, Neurosurgery & Psychiatry* 1990; 53(4): 293-296.
- Lundin AD, Boussard C, Edman G, Borg J. Symptoms and disability until 3 months after mild TBI. *Brain Injury* 2006; 20(8): 799-786.
- Maas, A.I., Menon, D.K., Lingsma, H.F., Pineda, J.A., Sandel, M.E. & Manley, G.T. Re-orientation of clinical research in traumatic brain injury: report of an international workshop on comparative effectiveness research. *J Neurotrauma* 2012; 29(1): 32-46.
- Mannix R, Meehan WP, Monuteaux MC, & Bachur RG. Computed tomography for minor head injury: variation and trends in major United States pediatric emergency departments. *The Journal of Pediatrics* 2012; 160(1): 136-139.
- Reuben A, Sampson P, Harris AR, Williams H, & Yates P. Post-concussion syndrome (PCS) in the emergency department: predicting and pre-empting persistent symptoms following a mild traumatic brain injury. *Emergency Medicine Journal: EMJ* 2014; 31(1): 72-77.
- Sigurdardottir S, Andelic N, Roe C, Jerstad T, & Schanke AK. Post-concussion symptoms after traumatic brain injury at 3 and 12 months post-injury: A prospective study. *Brain Injury* 2009; 23(6): 489-497.
- Stulemeijer M, van der Werf S, Borm GF, & Vos PE. Early prediction of favourable recovery 6 months after mild traumatic brain injury. *Journal Of Neurology, Neurosurgery, And Psychiatry* 2007; 79(8): 936-942.
- Thomson Microdex Health Care Service. (2002). Blunt head trauma. Retrieved February 24, 2013, from <http://www.headinjury.com/emergency.htm>



- Williams WH, Potter S, & Ryland H. Mild traumatic brain injury and Post-concussion Syndrome: a neuropsychological perspective. *Journal of Neurology, Neurosurgery & Psychiatry*. 2010; 81(10); 1116-1122.
- Wood RL. Understanding the miserable minority: A diathesis-stress paradigm for post-concussional syndrome. *Brain Injury* 2009; 18(11): 1135-1153.
- Wrightson, P., & Gronwall, D. *Mild Head Injury: A Guide to Management*. Oxford University Press; 1999
- Yang CC, Tu YK, Hua MS, & Huang SJ. The association between the postconcussion symptoms and clinical outcomes for patients with mild traumatic brain injury. *The Journal of Trauma* 2007; 62(3): 657-663.
- Zemek RL, Farion KJ, Sampson M, & McGahern C. Prognosticators of persistent symptoms following pediatric concussion: a systematic review. *JAMA Pediatrics* 2013;167(3): 259-265.